arXiv:math/0611501v3 [math.QA] 10 Aug 2007

VARIETIES OF DIALGEBRAS AND CONFORMAL ALGEBRAS

PAVEL KOLESNIKOV

Abstract. For a given variety Var of algebras we define the variety Var of dialgebras. This construction turns to be closely related with varieties of pseudo-algebras: every Var-dialgebra can be embedded into an appropriate pseudo-algebra of the variety Var. In particular, Leibniz algebras are exactly Lie dialgebras, and every Leibniz algebra can be embedded into current Lie conformal algebra.

Introduction

Leibniz algebras, introduced by J.-L. Loday [1], are non-commutative analogues of Lie algebras. The defining identity of the variety of left (right) Leibniz algebras is the condition stating that each operator of left (right) multiplication is a derivation, i.e.,

$$x(yz) = (xy)z + y(xz) \quad \text{or} \quad (xy)z = (xz)y + x(yz),$$

respectively. As Lie algebras are embeddable into associative algebras, Leibniz algebras can be embedded into associative dialgebras. By the definition [2, 3], dialgebras are linear spaces with two bilinear products $\dashv, \vdash$. The identities required for a dialgebra A to be associative are chosen in such a way that the new operation

$$ab = a \vdash b - b \dashv a \quad \text{or} \quad ab = a \dashv b - b \vdash a$$

turns A into a left (right) Leibniz algebra:

$$(x \dashv y) \vdash z = (x \vdash y) \vdash z, \quad x \dashv (y \vdash z) = x \dashv (y \dashv z), \tag{1}$$

$$(x \vdash y) \vdash z = x \vdash (y \vdash z), \quad (x \dashv y) \dashv z = x \dashv (y \dashv z), \tag{2}$$

$$(x \vdash y) \dashv z = x \vdash (y \dashv z). \tag{3}$$

In the paper [4] the notion of an alternative dialgebra was introduced. It was noted in [5] that an algebra A is embedded into the Steinberg Lie algebra $\mathrm{st}(n, A)$, $n \geq 4$, if and only if A is associative; and if $n = 3$ then A has to be at least alternative. In [4], the Steinberg Leibniz algebra $\mathrm{stl}(n, D)$ was introduced, where D is a dialgebra. The identities that are necessary for embedding of D into $\mathrm{stl}(3, D)$ are declared to be the identities of an alternative dialgebra.

The main purpose of this paper is to present a general scheme how to deduce what is a variety of dialgebras similar to a given variety of ordinary algebras (associative, Lie, alternative, etc.). This structure is motivated by the relation of dialgebras and conformal algebras.

A conformal algebra [6] is a linear space C (over a field of zero charachteristic) with one linear operation $T : C \to C$ and a countable family of bilinear products

---

Key words and phrases. Leibniz algebra, dialgebra, conformal algebra, operad, pseudo-algebra.

Partially supported by RFBR 05–01–00230, Complex Integration Program SB RAS (2006–1.9). The author gratefully acknowledges the support from Deligne 2004 Balzan prize in mathematics.

$(\cdot\ _{(n)}\ \cdot) : C \times C \to C$, where n ranges over the set $\mathbb{Z}_+$ of non-negative integers, such that

- for every $a, b \in C$ only a finite number of $a\ _{(n)}\ b$ is nonzero;
- $Ta\ _{(n)}\ b = -na\ _{(n-1)}\ b$, $a\ _{(n)}\ Tb = T(a\ _{(n)}\ b) + na\ _{(n-1)}\ b$.

Every conformal algebra can be constructed as a subspace of the formal power series space $A[[z, z^{-1}]]$ over an appropriate ordinary algebra A, with respect to

$$(Ta)(z) = \frac{d}{dz}a(z), \quad (a\ _{(n)}\ b)(z) = \operatorname*{Res}_{w=0} a(w)b(z)(w-z)^n.$$

Here $\operatorname{Res}_{w=0} f(w, z)$ stands for the coefficient of w^{-1} in $f(z, w) \in A[[z, z^{-1}, w, w^{-1}]]$.

In [7] it was shown that for an arbitrary conformal algebra C there exists unique ordinary algebra $\operatorname{Coeff} C$ (called coefficient algebra) which is universal in the following sense: $\operatorname{Coeff} C[[z, z^{-1}]]$ contains C, and for any algebra A such that $A[[z, z^{-1}]]$ contains C as above there exists a homomorphism $\operatorname{Coeff} C \to A$ such that the natural expansion $\operatorname{Coeff} C[[z, z^{-1}]] \to A[[z, z^{-1}]]$ acts on C as an identity.

A conformal algebra C is said to be associative (Lie, alternative, etc.) [7] if so is $\operatorname{Coeff} C$. A more general approach was developed in [8], see Section 1.4 for more details. For example, C is associative if

$$a\ _{(n)}\ (b\ _{(m)}\ c) = \sum_{s\geq 0} \binom{n}{s} (a\ _{(n-s)}\ b)\ _{(m+s)}\ c, \quad a, b, c \in C,\ n, m \in \mathbb{Z}_+.$$

It is not difficult to note that an associative conformal algebra C endowed with two operations

$$a \vdash b = a\ _{(0)}\ b, \quad a \dashv b = \sum_{s\geq 0} \frac{1}{s!}(-T)^s (a\ _{(s)}\ b)$$

satisfies (1)–(3), i.e., this is an associative dialgebra. The same is true for alternative conformal algebras. However, it is not obvious why every associative (or alternative) dialgebra can be obtained in this way. This is why we define what is a variety of dialgebras using the language of operads, and then prove that every dialgebra from a variety Var can be embedded into a conformal algebra (more generally, pseudo-algebra) from the variety Var of conformal algebras.

1. Operads and algebras

1.1. **Partitions.** Suppose $m \geq n \geq 1$ are two integers. An ordered n-tuple of integers $\pi = (m_1, \dots, m_n)$, $m_i \geq 1$, is called an *n-partition* of m if $m_1 + \dots + m_n = m$. The set of all such partitions is denoted by $\Pi(m, n)$.

A partition $\pi \in \Pi(m, n)$ determines a one-to-one correspondence between the sets $\{1, \dots, m\}$ and $\{(i, j) \mid i = 1, \dots, n,\ j = 1, \dots, m_i\}$ by the rule

$$(i, j) \leftrightarrow k = (i, j)^\pi, \quad k = m_1 + \dots + m_{i-1} + j.$$

Given two partitions $\tau = (p_1, \dots, p_m) \in \Pi(p, m)$, $\pi = (m_1, \dots, m_n) \in \Pi(m, n)$, define $\tau\pi \in \Pi(p, n)$ in a natural way:

$$\tau\pi = (p_1 + \dots + p_{m_1}, p_{m_1+1} + \dots + p_{m_1+m_2}, \dots, p_{m-m_n+1} + \dots + p_m).$$

If $\tau\pi = (q_1, \dots, q_n)$ then for any $i = 1, \dots, n$ denote by $\tau\pi_i$ the subpartition $(p_{m_1+\dots+m_{i-1}+1}, \dots, p_{m_1+\dots+m_i}) \in \Pi(q_i, m_i)$.

We will also consider the (right) action of the symmetric group S_n on $\Pi(m, n)$: if $\pi = (m_1, \dots, m_n)$, $\sigma \in S_n$ then $\pi\sigma = (m_{1\sigma^{-1}}, \dots, m_{n\sigma^{-1}})$.

1.2. **Operads.** Consider the following collection of data:

1) a family of linear spaces $\{\mathcal{C}(n)\}_{n\ge 1}$;
2) a family of linear maps (called composition rule)

$$\mathrm{Comp}^{\pi} : \mathcal{C}(n)\otimes\mathcal{C}(m_1)\otimes\cdots\otimes\mathcal{C}(m_n)\to\mathcal{C}(m),$$

where $\pi=(m_1,\dots,m_n)\in\Pi(m,n)$.

This collection is called an *operad* (denoted by $\mathcal{C}$) if the following properties hold.

(O1) The composition rule is associative, i.e., if $\tau=(p_1,\dots,p_m)\in\Pi(p,m)$, $\pi=(m_1,\dots,m_n)\in\Pi(m,n)$, and

$$\psi_j\in\mathcal{C}(p_j),\ \chi_i\in\mathcal{C}(m_i),\ \varphi\in\mathcal{C}(n),\quad i=1,\dots,n,\ j=1,\dots,m,$$

then

$$\mathrm{Comp}^{\tau}\big(\mathrm{Comp}^{\pi}\big(\varphi,(\chi_i)_{i=1}^{n}\big),(\psi_j)_{j=1}^{m}\big) = \mathrm{Comp}^{\tau\pi}\big(\varphi,\big(\mathrm{Comp}^{\tau\pi_i}\big(\chi_i,(\psi_{(i,t)^{\pi}})_{t=1}^{m_i}\big)_{i=1}^{n}\big)\big). \tag{4}$$

(O2) There exists a “unit” $\mathrm{id}\in\mathcal{C}(1)$ such that

$$\mathrm{Comp}^{(1,\dots,1)}(f,\mathrm{id},\dots,\mathrm{id})=\mathrm{Comp}^{(n)}(\mathrm{id}_A,f)=f \tag{5}$$

for every $f\in\mathcal{C}(n)$.

An important example of an operad is provided by the following data. Suppose $\mathcal{C}(n)=\Bbbk S_n$. If $\sigma\in S_n$, $\pi=(m_1,\dots,m_n)\in\Pi(m,n)$, $\tau_i\in S_{m_i}$, $i=1,\dots,n$, then define

$$\Sigma=\mathrm{Comp}^{\pi}(\sigma,\tau_1,\dots,\tau_n)\in S_m$$

by the rule: if $k=(i,j)^{\pi}\in\{1,\dots,m\}$ then $k\Sigma=(i\sigma,j\tau_i)^{\pi\sigma}$.

The principle of constructing such compositions is clearly shown by the following simple example:

$$\mathrm{Comp}^{(3,2,4)}((123),(132),(12),(234))=\begin{pmatrix}1&2&3&4&5&6&7&8&9\\7&5&6&9&8&1&3&4&2\end{pmatrix}.$$

Graphically, this composition can be expressed by the diagram



The operad obtained is called the *operad of symmetries* and denoted by Sym.

Suppose $\mathcal{C}$ is an operad. If $\mathcal{C}(n)$ is endowed with a right action of the symmetric group S_n for all $n\ge 1$, and for any $\pi=(m_1,\dots,m_n)\in\Pi(m,n)$, $\varphi\in\mathcal{C}(n)$, $\sigma\in S_n$, $\psi_i\in\mathcal{C}(m_i)$, $\tau_i\in S_{m_i}$, $i=1,\dots,n$, we have

$$\mathrm{Comp}^{\pi\sigma}\big(\varphi^{\sigma},\psi_{1\sigma^{-1}}^{\tau_{1\sigma^{-1}}},\dots,\psi_{n\sigma^{-1}}^{\tau_{n\sigma^{-1}}}\big)=\mathrm{Comp}^{\pi}(\varphi,\psi_1,\dots,\psi_n)^{\mathrm{Comp}^{\pi}(\sigma,\tau_1,\dots,\tau_n)},$$

then the operad $\mathcal{C}$ is said to be *symmetric*.

For example, Sym is a symmetric operad with respect to the regular right action of S_n on $\mathrm{Sym}(n)$: $\varphi^{\sigma}=\varphi\sigma$, $\sigma\in S_n$, $\varphi\in\mathrm{Sym}(n)=\Bbbk S_n$.

Let $\mathcal{C}_1$, $\mathcal{C}_2$ be two operads. A *functor* F from $\mathcal{C}_1$ to $\mathcal{C}_2$ is a family of linear maps $F_n : \mathcal{C}_1(n) \to \mathcal{C}_2(n)$, $n \geq 1$, which preserve the composition rule and the unit. If $\mathcal{C}_i$ are symmetric operads, and each F_n is S_n-invariant then F is said to be symmetric.

If $\mathcal{C}_1$, $\mathcal{C}_2$ are two (symmetric) operads, then the collection of spaces $\mathcal{C}(n) = \mathcal{C}_1(n) \otimes \mathcal{C}_2(n)$, $n \geq 1$, defines a (symmetric) operad with respect to the componentwise composition (and symmetric group action). We will denote this operad by $\mathcal{C}_1 \otimes \mathcal{C}_2$.

1.3. **Examples of operads.** We have already considered the operad Sym built on symmetric groups. Let us now state some well-known examples of operads that will be used later. Throughout the rest of the subsection, π stands for a partition $(m_1, \dots, m_n) \in \Pi(m, n)$, σ is a permutation from S_n.

1.3.1. *Operad* $\mathcal{E}$. Set $\mathcal{E}(n) = \Bbbk^n$, and let $\{e_1^{(n)}, \dots, e_n^{(n)}\}$ be the standard basis of $\Bbbk^n$. Define a composition rule and S_n-action as follows:

$$\mathrm{Comp}^\pi\big(e_i^{(n)}, e_{j_1}^{(m_1)}, \dots, e_{j_n}^{(m_n)}\big) = e^{(m)}_{m_1+\dots+m_{i-1}+j_i},$$

$$\sigma : e_i^{(n)} \mapsto e_{i\sigma}^{(n)}.$$

This is a symmetric operad. In fact, $\mathcal{E}(n)$ can be considered as an image of $\mathrm{Sym}(n)$ under the map $F_n : \Bbbk S_n \to \Bbbk^n$ given by $F_n(\sigma) = \sigma S_{n-1}$, assuming $(in)S_{n-1}$ is identified with $e_i^{(n)}$. The family of maps F_n, $n \geq 1$, defines a functor from Sym to $\mathcal{E}$, however, non-symmetric.

1.3.2. *Operads* $\mathrm{Vec}_\Bbbk$. Let V be a linear space over a field $\Bbbk$. Define an operad which is also denoted by V as follows:

$$V(n) = \mathrm{Hom}(V^{\otimes n}, V), \quad n \geq 1,$$

Comp is the ordinary composition of multi-linear maps. If $f \in V(n)$ then set

$$f^\sigma(v_1, \dots, v_n) = f(v_{1\sigma}, \dots, v_{n\sigma}), \quad v_i \in V.$$

This is a symmetric operad. We will denote the class of operads obtained in this way by $\mathrm{Vec}_\Bbbk$.

1.3.3. *Operads* H-mod. Suppose H is a cocommutative bialgebra with a coproduct $\Delta : H \to H \otimes H$. We will use the standard Sweedler's notation:

$$\Delta(h) = h_{(1)} \otimes h_{(2)}, \quad \Delta^2(h) = (\Delta \otimes \mathrm{id}_H)\Delta(h) = (\mathrm{id}_H \otimes \Delta)\Delta(h) = h_{(1)} \otimes h_{(2)} \otimes h_{(3)},$$

etc.

For every left H-module M one may consider

$$M(n) = \mathrm{Hom}_{H^{\otimes n}}(M^{\otimes n}, H^{\otimes n} \otimes_H M), \tag{6}$$

where $H^{\otimes n}$ is considered as the outer product of regular right H-modules.

An arbitrary $f \in M(n)$ can be expanded to a map

$$f : (H^{\otimes m_1} \otimes_H M) \otimes \dots \otimes (H^{\otimes m_n} \otimes_H M) \to H^{\otimes m} \otimes_H M$$

by the following rule: if $G_i \in H^{\otimes m_i}$, $a_i \in M$, $i = 1, \dots, n$, then

$$\begin{aligned} &f(G_1 \otimes_H a_1, \dots, G_n \otimes_H a_n) \\ &\quad = (G_1 \otimes \dots \otimes G_n \otimes_H 1)(\Delta^{m_1-1} \otimes \dots \otimes \Delta^{m_n-1} \otimes_H \mathrm{id}_M) f(a_1, \dots, a_n). \end{aligned} \tag{7}$$

This expansion makes clear what is a composition of maps (6). The action of S_n on $M(n)$ can be defined by

$$\varphi^\sigma(a_1,\dots,a_n) = (\sigma\otimes_H \mathrm{id}_M)\varphi(a_{1\sigma},\dots,a_{n\sigma}),$$

where $(h_1\otimes\dots\otimes h_n)^\sigma = (h_{1\sigma^{-1}}\otimes\dots\otimes h_{n\sigma^{-1}})\in H^{\otimes n}$.

We will denote the operad obtained in this way by the same symbol M. The class of all such operads is denoted by H-mod.

1.3.4. *Operad* Alg. Let $X=\{x_1,x_2,\dots\}$ be a countable set of symbols. Define $\mathrm{Alg}(n)$ to be the linear span of all non-associative words obtained from $x_1\dots x_n$ by some bracketing. The natural composition rule

$$\mathrm{Comp}^\pi(u,v_1,\dots,v_n) = u(v_1(x_{(1,1)^\pi},\dots,x_{(1,m_1)^\pi}),\dots,v_n(x_{(n,1)^\pi},\dots,x_{(n,m_n)^\pi})) \tag{8}$$

defines a (non-symmetric) operad denoted by Alg.

Consider $\mathrm{Alg}_S(n)=\mathrm{Alg}(n)\otimes \Bbbk S_n$ endowed with a composition rule

$$\begin{aligned}\mathrm{Comp}^\pi&(u\otimes\sigma, v_1\otimes\tau_1,\dots,v_n\otimes\tau_n)\\ &= \mathrm{Comp}^{\pi\sigma^{-1}}(u,v_{1\sigma},\dots,v_{n\sigma})\otimes \mathrm{Comp}^{\pi\sigma^{-1}}(\sigma,\tau_{1\sigma},\dots,\tau_{n\sigma}).\end{aligned} \tag{9}$$

Hereafter, we will identify $(x_1\dots x_n)\otimes\sigma$ with the word $(x_{1\sigma}\dots x_{n\sigma})$. Under the natural action of S_n, Alg_S is a symmetric operad.

If $\mathcal{C}$ is a symmetric operad and Φ is a functor from Alg to $\mathcal{C}$ then Φ can be extended to a functor from Alg_S to $\mathcal{C}$ by the obvious rule

$$\Phi_n(u\otimes\sigma)=\Phi_n(u)^\sigma,\quad u\in \mathrm{Alg}(n),\ \sigma\in S_n.$$

1.3.5. *Operad* VarAlg. Let Var be a homogeneous variety of (non-associative) algebras defined by a family of polylinear identities. Denote by $\Bbbk_{\mathrm{Var}}\{X\}$ the free algebra in this variety generated by $X=\{x_1,x_2,\dots\}$. There exists an ideal I_{Var} of the free algebra $\Bbbk\{X\}$ such that $\Bbbk_{\mathrm{Var}}\{X\}\simeq \Bbbk\{X\}/I_{\mathrm{Var}}$.

Each space $\mathrm{Alg}_S(n)$, $n\ge 1$, can be considered as a subspace of $\Bbbk\{X\}$. Denote

$$\mathrm{VarAlg}(n)=\mathrm{Alg}_S(n)/(\mathrm{Alg}_S(n)\cap I_{\mathrm{Var}}).$$

Since the variety Var is homogeneous, the composition map and the symmetric group action are well-defined on the family $\{\mathrm{VarAlg}(n)\}_{n\ge1}$, so we obtain a symmetric operad. The family of projections $\mathrm{Alg}_S(n)\to \mathrm{VarAlg}(n)$ is a functor from Alg_S to VarAlg, we will also denote it by Var.

1.4. **Algebras and pseudo-algebras.** Let A stands for a (non-associative, in general) algebra over $\Bbbk$. Then there exists a functor (also denoted by A) from Alg to the operad $A\in \mathrm{Vec}_\Bbbk$ built on the underlying space. Indeed, for every $u=(x_1\dots x_n)\in \mathrm{Alg}(n)$ define $A_n(u): a_1\otimes\dots\otimes a_n\to (a_1\dots a_n)\in A$.

Conversely, every functor from Alg to $A\in\mathrm{Vec}_\Bbbk$ defines an operation $A\otimes A\to A$ which makes A to be an algebra.

Therefore, the notion of an algebra can be identified with the notion of a functor from Alg to an operad from $\mathrm{Vec}_\Bbbk$. Similarly, a pseudo-algebra [9] is just a functor from Alg to an operad from H-mod. Since $\mathrm{Vec}_\Bbbk$ and H-mod consist of symmetric operads, a (pseudo-)algebra can be considered as a functor defined on $\mathrm{Alg}_S(n)$.

This is now clear how to define varieties of pseudo-algebras. Indeed, suppose Var is a variety as in 1.3.5. An (ordinary) algebra A belongs to Var if and only if there exists a functor $\bar{A} : \mathrm{VarAlg} \to A \in \mathrm{Vec}_{\Bbbk}$ such that $A = \mathrm{Var} \circ \bar{A}$. For pseudo-algebras, one should replace $\mathrm{Vec}_{\Bbbk}$ with H-mod to obtain a definition of what does it mean that a pseudo-algebra C belongs to the variety Var [10].

This approach leads to pseudo-algebraic identities as in [8]. If Var is defined by a family Σ of polylinear identities then a pseudo-algebra C is said to be a Var-pseudo-algebra if C satisfies the identities

$$t^*(x_1, \dots, x_n), \quad t \in \Sigma,$$

where t^* is obtained from t by the following way: every monomial $\alpha(x_{1\sigma} \dots x_{n\sigma})$, $\alpha \in \Bbbk$, $\sigma \in S_n$, that appears in t should be replaced with $\alpha(\sigma \otimes_H \mathrm{id})(x_{1\sigma} * \dots * x_{n\sigma})$. The expression $(x_{1\sigma} * \dots * x_{n\sigma})$ can be evaluated at $a_1, \dots, a_n \in C$ by (7), its value lies in $H^{\otimes n} \otimes_H C$.

2. Dialgebras and corresponding operads

Consider an operad Dialg, analogous to Alg, generated by two binary products $\dashv, \vdash$. The space $\mathrm{Dialg}(n)$ is the linear span of all terms obtained from $x_1 \dots x_n$ by some bracketing with operations $\dashv, \vdash$, and the same rule as (8) defines compositions on Dialg.

By Dialg_S we denote the symmetric operad obtained from Dialg in the same way as Alg_S is obtained from Alg. The operad Dialg (resp., Dialg_S) has the following property: for any (symmetric) operad $\mathcal{C}$ and for any $\varphi_1, \varphi_2 \in \mathcal{C}(2)$ there exists a unique (symmetric) functor $\Psi : \mathrm{Dialg} \to \mathcal{C}$ ($\mathrm{Dialg}_S \to \mathcal{C}$) such that $\Psi_2(x_1 \vdash x_2) = \varphi_1$, $\Psi_2(x_1 \dashv x_2) = \varphi_2$. In particular, for $\mathcal{C} = \mathrm{Alg}_S \otimes \mathcal{E}$ there exists $\Psi : \mathrm{Dialg}_S \to \mathrm{Alg}_S \otimes \mathcal{E}$ such that

$$\Psi_2(x_1 \vdash x_2) = x_1 x_2 \otimes e_2^{(2)}, \quad \Psi_2(x_1 \dashv x_2) = x_1 x_2 \otimes e_1^{(2)}. \tag{10}$$

It is clear that the functor Ψ defined by (10) is full, i.e., $\Psi_n(\mathrm{Dialg}_S(n)) = \mathrm{Alg}_S(n) \otimes \mathcal{E}(n)$. Throughout the rest of the paper, Ψ stands for the functor $\mathrm{Dialg}_S \to \mathrm{Alg}_S \otimes \mathcal{E}$ defined by (10).

Definition 1. A dialgebra A is said to be a 0-*dialgebra* if A satisfies the identities

$$(x \dashv y) \vdash z = (x \vdash y) \vdash z, \quad x \dashv (y \vdash z) = x \dashv (y \dashv z). \tag{11}$$

The purpose of this section is to prove the following statement.

Theorem 1. *For any 0-dialgebra A, considered as a functor $\mathrm{Dialg}_S \to A \in \mathrm{Vec}_{\Bbbk}$, there exists a symmetric functor $\bar{A} : \mathrm{Alg}_S \otimes \mathcal{E} \to A$ such that $A = \Psi \circ \bar{A}$.*

Let H be a cocommutative Hopf algebra with a counit ε and an antipode S.

If C is a left H-module then $H^{\otimes n} \otimes_H C \simeq H^{\otimes n-1} \otimes C$ as a linear space [9]. We will denote the isomorphism by i_n ($n \geq 1$), namely,

$$i_n(h_1 \otimes \dots \otimes h_n \otimes_H a) = h_1 S(h_{n(1)}) \otimes \dots \otimes h_{n-1} S(h_{n(n-1)}) \otimes h_{n(n)} a,$$

$h_i \in H$, $a \in C$.

For an operad $C \in H$-mod denote by $C^{(0)}$ the operad from $\mathrm{Vec}_{\Bbbk}$ built on C as on linear space.

Lemma 2. *A family of linear maps*

$$\varepsilon_n : C(n) \to C^{(0)}(n), \quad n \geq 1,$$

given by

$$\varepsilon_n(f)(a_1,\dots,a_n) = (\varepsilon^{\otimes n-1}\otimes \mathrm{id})i_n(f(a_1,\dots,a_n)), \quad f\in C(n),$$

defines a (non-symmetric) functor from C *to* $C^{(0)}$.

Proof. The expansion rule (7) immediately implies that for all $f\in C(n)$, $g_i\in C(m_i)$, $i=1,\dots,n$, $\pi=(m_1,\dots,m_n)\in\Pi(m,n)$, we have

$$\varepsilon_m(\mathrm{Comp}^\pi(f,g_1,\dots,g_n)) = \mathrm{Comp}^\pi(\varepsilon_n(f),\varepsilon_{m_1}(g_1),\dots,\varepsilon_{m_n}(g_n)).$$

Therefore, the family of maps ε_n, $n\ge 1$, defines a functor (also denoted by ε). $\square$

Let C stands for an H-pseudo-algebra considered as a functor $\mathrm{Alg}_S\to C\in H\text{-mod}$. By the universal property of Dialg_S we have a uniquely defined symmetric functor from Dialg_S to $C^{(0)}\in \mathrm{Vec}_\Bbbk$ (also denoted by $C^{(0)}$) such that

$$\begin{gathered} C_2^{(0)}(x_1\vdash x_2) = \varepsilon_2(C_2(x_1x_2))\in C^{(0)}(2),\\ C_2^{(0)}(x_1\dashv x_2) = \varepsilon_2(C_2(x_2x_1))^{(12)}\in C^{(0)}(2). \end{gathered}\tag{12}$$

In particular, for every $a,b\in C$

$$\begin{gathered} a\vdash b = C_2^{(0)}(x_1\vdash x_2)(a,b) = (\varepsilon\otimes\mathrm{id})i_2(a*b),\\ a\dashv b = C_2^{(0)}(x_1\dashv x_2)(a,b) = \varepsilon_2(C_2(x_2x_1))(b,a) = (\varepsilon\otimes\mathrm{id})i_2\{a*b\}, \end{gathered}$$

where $\{a*b\} = ((12)\otimes_H \mathrm{id})(a*b)$.

Lemma 3. *For every H-pseudo-algebra C the corresponding dialgebra $C^{(0)}$ given by* (12) *is a 0-dialgebra.*

Proof. It is straightforward to check that the identities (11) hold on $C^{(0)}$. $\square$

Proposition 4. *For every 0-dialgebra A there exists an H-pseudo-algebra $C=C(A)$ over $H=\Bbbk[T]$ such that A is embedded into $C^{(0)}$.*

Proof. Here we consider $\Bbbk[T]$ with respect to the usual Hopf algebra structure: $\Delta(T)=T\otimes 1+1\otimes T$, $\varepsilon(T)=0$, $S(T)=-T$.

Let A be a 0-dialgebra. Denote $\langle a,b\rangle = a\vdash b - a\dashv b$. It follows from (11) that $\langle a,b\rangle\vdash c=0$, $a\dashv\langle b,c\rangle=0$ for all $a,b,c\in A$.

Consider the linear space

$$C=(H\otimes A)\oplus\left((A\otimes A)/\operatorname{Span}\{\langle a,b\rangle\otimes\langle c,d\rangle\mid a,b,c,d\in A\}\right).$$

The first summand is a free H-module, the second one can be endowed with an H-module structure as follows. Define $T: A\otimes A\to A$ by

$$T(a\otimes b)=\langle a,b\rangle,\quad a,b\in A,\tag{13}$$

and note that $U=\operatorname{Span}\{\langle a,b\rangle\otimes\langle c,d\rangle\mid a,b,c,d\in A\}$ lies in the kernel of T. Therefore, T is well-defined on $(A\otimes A)/U$.

Later we will identify $1\otimes a\in H\otimes A$ with a, and write $a\otimes b$ ($a,b\in A$) for its image in C.

The pseudo-product on C can be defined as follows:

$$a*b = 1\otimes 1\otimes_H (a\vdash b) - T\otimes 1\otimes_H (a\otimes b),\tag{14}$$

$$a*(b\otimes c) = 1\otimes 1\otimes_H a\otimes\langle b,c\rangle = 1\otimes 1\otimes_H a\otimes T(b\otimes c),\tag{15}$$

$$(a\otimes b)*c = -1\otimes 1\otimes_H (\langle a,b\rangle\otimes c) = -1\otimes 1\otimes_H (T(a\otimes b)\otimes c),\tag{16}$$

$$(a\otimes b)*(c\otimes d)=0.\tag{17}$$

This operation is well-defined and $H^{\otimes 2}$-linear. It is easy to see that A is a subdialgebra of $C^{(0)}$. $\square$

Lemma 5. *Suppose C is an H-pseudo-algebra. Consider a family of linear maps*

$$C_n^{(\varepsilon)} : \mathrm{Alg}_S(n) \otimes \mathcal{E}(n) \to C^{(0)}(n), \quad n \geq 1,$$

defined by the following rule: if $C_n(f)(a_1, \dots, a_n) = \sum\limits_j h_{1j} \otimes \dots \otimes h_{nj} \otimes_H b_j$, $f \in \mathrm{Alg}_S(n)$, $a_i, b_j \in C$, $h_{ij} \in H$, then

$$C_n^{(\varepsilon)}(f \otimes e_i)(a_1, \dots, a_n) = \sum_j \varepsilon(h_{1j} \overset{i}{\widehat{\ldots}} h_{nj}) h_{ij} b_j.$$

These maps define a symmetric functor $C^{(\varepsilon)} : \mathrm{Alg}_S \otimes \mathcal{E} \to C^{(0)} \in \mathrm{Vec}_\Bbbk$.

Proof. This is straightforward to check that

$$\big(C_n^{(\varepsilon)}\big(f \otimes e_i^{(n)}\big)\big)^\sigma = C_n^{(\varepsilon)}\big(f^\sigma \otimes e_{i\sigma}^{(n)}\big), \quad f \in \mathrm{Alg}_S(n),\ i = 1, \dots, n,\ \sigma \in S_n.$$

It remains to prove that $C_n^{(\varepsilon)}$ preserve compositions. Since the composition rule is equivariant and associative, it is enough to consider $\pi = (n, m-n) \in \Pi(m, 2)$, $x_1x_2 \in \mathrm{Alg}(2)$, $u_1 \in \mathrm{Alg}(n)$, $u_2 \in \mathrm{Alg}(m-n)$, and

$$\begin{aligned} C_m^{(\varepsilon)}\big(\mathrm{Comp}^\pi\big(x_1x_2 \otimes e_i^{(2)}, u_1 \otimes e_k^{(n)}, u_2 \otimes e_l^{(m-n)}\big)\big) \\ = C_m^{(\varepsilon)}\big(f \otimes e_p^{(m)}\big), \quad p = \begin{cases} k, & i = 1, \\ n + l, & i = 2, \end{cases} \end{aligned} \tag{18}$$

where $f = u_1(x_1, \dots, x_n) u_2(x_{n+1}, \dots, x_m)$. Now it is easy to note that

$$\mathrm{Comp}^{(n,m-n)}\big(C_2^{(\varepsilon)}\big(x_1x_2 \otimes e_i^{(2)}\big), C_n^{(\varepsilon)}\big(u_1 \otimes e_k^{(n)}\big), C_{m-n}^{(\varepsilon)}\big(u_2 \otimes e_l^{(m-n)}\big)\big)$$

coincides with (18). $\square$

Proof of Theorem 1. Let A be a 0-dialgebra. Denote by C its enveloping pseudo-algebra from Proposition 4. The functor $C : \mathrm{Alg}_S \to C \in H\text{-mod}$ induces two functors $C^{(0)} : \mathrm{Dialg}_S \to C^{(0)}$, $C^{(\varepsilon)} : \mathrm{Alg}_S \otimes \mathcal{E} \to C^{(0)}$. It is easy to see that $\Psi \circ C^{(\varepsilon)} = C^{(0)}$. Therefore, if $\Psi_n(f) = 0$ for some $f \in \mathrm{Dialg}_S(n)$ then $C_n^{(0)}(f) = 0$ in $C^{(0)}(n)$. But A is embedded into $C^{(0)}$, hence, f is an identity on A_0.

Therefore, a polynomial $f \in \mathrm{Dialg}_S(n)$ is an identity on a 0-dialgebra A provided that $\Psi_n(f) = 0$. Hence, there exists a family of linear maps $\bar{A}_n : \mathrm{Alg}_S(n) \otimes \mathcal{E}(n) \to A(n)$ such that $A_n = \Psi_n \otimes \bar{A}_n$, $n \geq 1$. This family defines a symmetric functor $\bar{A}$ from $\mathrm{Alg}_S \otimes \mathcal{E}$ to $A \in \mathrm{Vec}_\Bbbk$. $\square$

One may construct an operad 0-Dialg related to the variety of 0-dialgebras in the same way as VarAlg relates to a variety Var.

Corollary 6. *The operad 0-*Dialg *is equivalent to* $\mathrm{Alg}_S \otimes \mathcal{E}$.

3. VARIETIES OF DIALGEBRAS

Suppose Var is a homogeneous variety defined by a family Σ of polylinear identities in $X = \{x_1, x_2, \dots\}$. Then I_{Var} is an ideal of $\Bbbk\{X\}$ generated by $\{t(f_1, \dots, f_n) \mid t(x_1, \dots, x_n) \in \Sigma,\ f_i \in \Bbbk\{X\}\}$.

Theorem 1 implies that every 0-dialgebra A is actually a functor from $\mathrm{Alg}_S \otimes \mathcal{E}$ to $A \in \mathrm{Vec}_\Bbbk$. In view of this observation, the following definition is quite natural.

Definition 2. A dialgebra $A : \mathrm{Dialg}_S \to A \in \mathrm{Vec}_\Bbbk$ is said to be a member of the variety Var of dialgebras (or Var-*dialgebra*) if there exists a symmetric functor

$$\bar{A} : \mathrm{VarAlg} \otimes \mathcal{E} \to A \in \mathrm{Vec}_\Bbbk,$$

such that $\Psi \circ (\mathrm{Var} \otimes \mathrm{id}) \circ \bar{A} = A$, i.e., the diagram of functors

$$\begin{tikzcd}
\mathrm{Dialg}_S \arrow[r, "A"] \arrow[d, "\Psi"'] & A \in \mathrm{Vec}_\Bbbk \\
\mathrm{Alg}_S \otimes \mathcal{E} \arrow[r, "\mathrm{Var}\otimes\mathrm{id}"] & \mathrm{VarAlg} \otimes \mathcal{E} \arrow[u, "\bar{A}"']
\end{tikzcd}$$

is commutative.

Proposition 7. *A dialgebra A belongs to the variety* Var *if and only if A is a 0-dialgebra that satisfies the identities $\Psi_n^{-1}\big(t \otimes e_i^{(n)}\big)$, where $t \in \Sigma$, $n = \deg t$, $i = 1, \dots, n$.*

Proof. It is enough to prove the "if" part only. Since A is a 0-dialgebra, it can be considered as a functor $A : \mathrm{Alg}_S \otimes \mathcal{E} \to A \in \mathrm{Vec}_\Bbbk$.

Suppose $f \in (\mathrm{Alg}_S \otimes \mathcal{E})(n)$, $f = \sum\limits_{i=1}^{n} f_i \otimes e_i^{(n)}$, $f_i \in \mathrm{Alg}_S(n)$. If $(\mathrm{Var}_n \otimes \mathrm{id})(f) = 0$ then $\mathrm{Var}_n(f_i) = 0$ for all $i = 1, \dots, n$. Therefore, it needs to be shown that if $\mathrm{Var}_n(g) = 0$ for some $g \in \mathrm{Alg}_S(n)$ then $A_n\big(g \otimes e_i^{(n)}\big) = 0$ for all $i = 1, \dots, n$. Since $\mathrm{Ker}\,\mathrm{Var}_n = \mathrm{Alg}_S(n) \cap I_{\mathrm{Var}}$, the element g can be presented as

$$g = \sum_k \mathrm{Comp}^{\pi_k}(a_k, b_{k1}, \dots, b_{kl_k})^{\sigma_k},$$

where $a_k \in \mathrm{Alg}_S(l_k)$, $b_{kj} \in \mathrm{Alg}_S(m_{kj})$, $\pi_k = (m_{k1}, \dots, m_{kl_k}) \in \Pi(n, l_k)$, $\sigma_k \in S_n$, and for each k there exists $j_k \in \{1, \dots, l_k\}$ such that

$$b_{kj_k} = \mathrm{Comp}^{\tau_k}(t_k, c_{k1}, \dots, c_{km_k}), \quad t_k \in \Sigma, \quad c_{km} \in \mathrm{Alg}_S(q_m).$$

For every $i \in \{1, \dots, n\}$ one may present $g \otimes e_i^{(n)}$ as a composition in $\mathrm{Alg}_S \otimes \mathcal{E}$ with the same a_k, b_{kj}, t_k, c_{km} in the first tensor factor (since the composition rule in $\mathrm{Alg}_S \otimes \mathcal{E}$ is componentwise). Therefore, $A_n\big(g \otimes e_i^{(n)}\big) = 0$ provided that $A_{m_k}\big(t_k \otimes e_p^{(m_k)}\big) = 0$ for every $p = 1, \dots, m_k$. $\square$

To deduce defining identities in the signature of a dialgebra, the following construction is useful. Consider the functor $\alpha : \mathrm{Dialg} \to \mathrm{Alg} \otimes \mathrm{Sym}$ defined by $\alpha_2(x_1 \vdash x_2) = x_1x_2 \otimes \mathrm{id}_2$, $\alpha_2(x_1 \dashv x_2) = x_1x_2 \otimes (12)$. Denote by E the following functor from Sym to $\mathcal{E}$: $E_n(\sigma) = e^{(n)}_{n\sigma^{-1}}$.

Lemma 8. *If $u \otimes \sigma \in \mathrm{Dialg}_S(n)$, $\alpha_n(u) = v \otimes \tau$, then*

$$\Psi_n(u \otimes \sigma) = (v \otimes \sigma) \otimes E_n(\tau)^\sigma. \tag{19}$$

Proof. Denote the right-hand side of (19) by $\Phi_n(u \otimes \sigma)$. It is enough to check that

1) $\Phi_n((u \otimes \sigma))^{\sigma_1} = \Phi_n(u \otimes \sigma\sigma_1)$;

2) the maps Φ_n, $n \geq 1$, preserve composition.

The first is obvious. The second needs to be checked for $u = u_1 \vdash u_2$ and $u = u_1 \dashv u_2$, $\sigma = \mathrm{id}_n$, $u_i \in \mathrm{Dialg}(n_i)$, $n_1 + n_2 = n$.

If $u = u_1 \vdash u_2$, $\alpha_{n_i}(u_i) = v_i \otimes \tau_i$, then $\alpha_n(u) = v_1v_2 \otimes \mathrm{Comp}^{(n_1,n_2)}(\mathrm{id}_2, \tau_1, \tau_2)$. Hence,

$$\Phi_n(u \otimes \mathrm{id}_n) = v_1v_2 \otimes \mathrm{id}_n \otimes e^{(n)}_{n_1+n_2\tau_2^{-1}}.$$

On the other hand,

$$\begin{gathered}\mathrm{Comp}^{(n_1,n_2)}\big(x_1x_2 \otimes e^{(2)}_2, \Phi_{n_1}(v_1 \otimes \mathrm{id}_{n_1}), \Phi_{n_2}(v_2 \otimes \mathrm{id}_{n_2})\big)\\ = (v_1v_2 \otimes \mathrm{id}_n) \otimes \mathrm{Comp}^{(n_1,n_2)}\big(e^{(2)}_2, e^{(n_1)}_{n_1\tau_1^{-1}}, e^{(n_2)}_{n_2\tau_2^{-1}}\big) = v_1v_2 \otimes \mathrm{id}_n \otimes e^{(n)}_{n_1+n_2\tau_2^{-1}}.\end{gathered}$$

If $u = u_1 \dashv u_2$, $\alpha_{n_i}(u_i) = v_i \otimes \tau_i$, then $\alpha_n(u) = v_1v_2 \otimes \mathrm{Comp}^{(n_1,n_2)}((12), \tau_1, \tau_2)$. Hence,

$$\Phi_n(u \otimes \mathrm{id}_n) = v_1v_2 \otimes \mathrm{id}_n \otimes e^{(n)}_{n_1\tau_1^{-1}}.$$

On the other hand,

$$\begin{gathered}\mathrm{Comp}^{(n_1,n_2)}\big(x_1x_2 \otimes e^{(2)}_1, \Phi_{n_1}(v_1 \otimes \mathrm{id}_{n_1}), \Phi_{n_2}(v_2 \otimes \mathrm{id}_{n_2})\big)\\ = (v_1v_2 \otimes \mathrm{id}_n) \otimes \mathrm{Comp}^{(n_1,n_2)}\big(e^{(2)}_1, e^{(n_1)}_{n_1\tau_1^{-1}}, e^{(n_2)}_{n_2\tau_2^{-1}}\big) = v_1v_2 \otimes \mathrm{id}_n \otimes e^{(n)}_{n_1\tau_1^{-1}}.\end{gathered}$$

Therefore, the sequence Φ_n, $n \geq 1$, defines a symmetric functor Φ from Dialg_S to $\mathrm{Alg}_S \otimes \mathcal{E}$ that coincides with Ψ on $\mathrm{Dialg}_S(2)$. Hence, $\Phi = \Psi$. $\square$

In order to get $\Psi_n^{-1}\big(v\otimes\sigma\otimes e^{(n)}_i\big)$ for some word $v \in \mathrm{Alg}(n)$, $\sigma \in S_n$, $i \in \{1, \dots, n\}$, it is enough to put the signs $\dashv$, $\vdash$ of dialgebra operations on the word v with the same bracketing in such a way that for the dialgebraic word $v^d_i \in \mathrm{Dialg}(n)$ obtained one has $\alpha_n\big(v^d_i\big) = v \otimes \tau$, where $i\sigma^{-1} = n\tau^{-1}$. Then $v^d_i \otimes \sigma \in \mathrm{Dialg}_S(n)$ would be a preimage of $v \otimes \sigma \otimes e^{(n)}_i$ with respect to Ψ_n.

The method is quite clear. Suppose $v \otimes \sigma = (x_{1\sigma} \dots x_{n\sigma}) \in \mathrm{Alg}_S(n)$. Then $v^d_i \otimes \sigma = (x_{1\sigma} \vdash \dots \vdash x_i \dashv \dots \dashv x_{n\sigma})$, i.e., the dashes are always directed to the variable x_i.

Let us consider some examples of varieties.

Associative dialgebras. In this case $\Sigma = \{(x_1, x_2, x_3)\}$, where $(x, y, z) = (xy)z - x(yz)$. Lemma 8 implies that

$$\begin{aligned}\Psi_3^{-1}\big((x_1, x_2, x_3) \otimes e^{(3)}_1\big) &= (x_1, x_2, x_3)_{\dashv},\\ \Psi_3^{-1}\big((x_1, x_2, x_3) \otimes e^{(3)}_2\big) &= (x_1, x_2, x_3)_{\times},\\ \Psi_3^{-1}\big((x_1, x_2, x_3) \otimes e^{(3)}_3\big) &= (x_1, x_2, x_3)_{\vdash},\end{aligned} \tag{20}$$

where

$$\begin{gathered}(x, y, z)_{\dashv} = (x \dashv y) \dashv z - x \dashv (y \dashv z), \quad (x, y, z)_{\times} = (x \vdash y) \dashv z - x \vdash (y \dashv z),\\ (x, y, z)_{\vdash} = (x \vdash y) \vdash z - x \vdash (y \vdash z).\end{gathered}$$

Together with (11), the identities (20) are exactly those introduced in [2, 3].

Commutative dialgebras. Let $\Sigma = \{(x_1, x_2, x_3),\ x_1x_2 - x_2x_1\}$. The first identity implies (20) as before, the last one is equivalent to

$$x_1 \vdash x_2 - x_2 \dashv x_1. \tag{21}$$

Therefore, a commutative dialgebra A can be considered as an ordinary algebra with respect to $ab = a \vdash b$, $a, b \in A$. In this case, we obtain an associative algebra that satisfies

$$[x_1, x_2]x_3 = 0. \tag{22}$$

Conversely, an arbitrary associative algebra that satisfies (22) is actually a commutative dialgebra.

Alternative dialgebras. Suppose $\Sigma = \{(x_1, x_2, x_3) + (x_2, x_1, x_3),\ (x_1, x_2, x_3) + (x_1, x_3, x_2)\}$. We have already seen what is obtained from $(x_1, x_2, x_3) \otimes e_i^{(3)}$ under Ψ_3^{-1}. Therefore, the identities Σ give rise to

$$\begin{gathered} (x_1, x_2, x_3)_\dashv + (x_1, x_2, x_3)_\times^{(12)}, \quad (x_1, x_2, x_3)_\vdash + (x_1, x_2, x_3)_\vdash^{(12)}, \\ (x_1, x_2, x_3)_\dashv + (x_1, x_2, x_3)_\dashv^{(23)}, \quad (x_1, x_2, x_3)_\times + (x_1, x_2, x_3)_\vdash^{(23)}. \end{gathered} \tag{23}$$

These relations are equivalent to those introduced in [4].

Lie dialgebras. If $\Sigma = \{(x_1, x_2, x_3) - x_2(x_1x_3),\ x_1x_2 + x_2x_1\}$ then the corresponding dialgebra identities include

$$x_1 \dashv x_2 + x_2 \vdash x_1.$$

A Lie dialgebra A considered as an ordinary algebra with respect to $[ab] = a \vdash b$, $a, b \in A$, is just a left Leibniz algebra. Conversely, every left Leibniz algebra L is a Lie dialgebra with respect to $a \vdash b = [ab]$, $a \dashv b = -[ba]$. Therefore, a Lie dialgebra is just the same as a Leibniz algebra.

Jordan dialgebras. Let Σ consists of the commutativity identity and

$$\begin{aligned} &x_1(x_2(x_3x_4)) + (x_2(x_1x_3))x_4 + x_3(x_2(x_1x_4)) \\ &\qquad = (x_1x_2)(x_3x_4) + (x_1x_3)(x_2x_4) + (x_3x_2)(x_1x_4). \end{aligned} \tag{24}$$

The last one is a polylinear Jordan identity.

Therefore, a Jordan dialgebra satisfies commutativity relation $x_1 \dashv x_2 = x_2 \vdash x_1$ and the following four identities:

$$\begin{aligned} &x_1 \dashv (x_2 \dashv (x_3 \dashv x_4)) + (x_2 \vdash (x_1 \dashv x_3)) \dashv x_4 + x_3 \vdash (x_2 \vdash (x_1 \dashv x_4)) \\ &\qquad = (x_1 \dashv x_2) \dashv (x_3 \dashv x_4) + (x_1 \dashv x_3) \dashv (x_2 \dashv x_4) + (x_3 \vdash x_2) \vdash (x_1 \dashv x_4). \end{aligned}$$

$$\begin{aligned} &x_1 \vdash (x_2 \dashv (x_3 \dashv x_4)) + (x_2 \dashv (x_1 \dashv x_3)) \dashv x_4 + x_3 \vdash (x_2 \dashv (x_1 \dashv x_4)) \\ &\qquad = (x_1 \vdash x_2) \dashv (x_3 \dashv x_4) + (x_1 \vdash x_3) \vdash (x_2 \dashv x_4) + (x_3 \vdash x_2) \dashv (x_1 \dashv x_4). \end{aligned}$$

$$\begin{aligned} &x_1 \vdash (x_2 \vdash (x_3 \dashv x_4)) + (x_2 \vdash (x_1 \vdash x_3)) \dashv x_4 + x_3 \dashv (x_2 \dashv (x_1 \dashv x_4)) \\ &\qquad = (x_1 \vdash x_2) \vdash (x_3 \dashv x_4) + (x_1 \vdash x_3) \dashv (x_2 \dashv x_4) + (x_3 \dashv x_2) \dashv (x_1 \dashv x_4). \end{aligned}$$

$$\begin{aligned} &x_1 \vdash (x_2 \vdash (x_3 \vdash x_4)) + (x_2 \vdash (x_1 \vdash x_3)) \vdash x_4 + x_3 \vdash (x_2 \vdash (x_1 \vdash x_4)) \\ &\qquad = (x_1 \vdash x_2) \vdash (x_3 \vdash x_4) + (x_1 \vdash x_3) \vdash (x_2 \vdash x_4) + (x_3 \vdash x_2) \vdash (x_1 \vdash x_4). \end{aligned}$$

Being expressed in terms of single operation, say, $xy = x \vdash y$, these identities together with (11) turn into

$$(x_1x_2)x_3 = (x_2x_1)x_3. \tag{25}$$

$$\begin{aligned}((x_4x_3)x_2)x_1 + x_4(x_2(x_3x_1)) + x_3(x_2(x_4x_1)) \\ = (x_4x_3)(x_2x_1) + (x_4x_2)(x_3x_1) + (x_3x_2)(x_4x_1).\end{aligned} \tag{26}$$

$$\begin{aligned}x_1((x_4x_3)x_2) + x_4((x_3x_1)x_2) + x_3((x_4x_1)x_2) \\ = (x_4x_3)(x_1x_2) + (x_1x_3)(x_4x_2) + (x_4x_1)(x_3x_2).\end{aligned} \tag{27}$$

Identities (26), (27) and (25) define a variety of algebras that relate to Jordan algebras as Leibniz algebras relate to Lie algebras. These algebras inherit the important property of Jordan algebras: the commutator of two operators of left multiplication is a derivation of the algebra.

4. Relation to pseudo-algebras

Suppose Var is a homogeneous variety defined by a family of polylinear identities Σ.

Theorem 9. *If C is an* Var*-pseudo-algebra over a cocommutative Hopf algebra H then $C^{(0)}$ is a* Var*-dialgebra.*

Proof. Every identity $t \in \Sigma$ of degree n can be considered as an element of $\mathrm{Alg}_S(n)$, i.e., $t = \sum\limits_{\sigma\in S_n} t_\sigma \otimes \sigma$, $t_\sigma \in \mathrm{Alg}(n)$.

Let us fix a linear basis $\{h_i\}$ of H and some elements $a_1, \dots, a_n \in C$. Denote

$$C_n(t_\sigma \otimes \sigma)(a_1, \dots, a_n) = \sum_{i_1,\dots,i_{n-1}} h_{i_1} \otimes \dots \otimes h_{i_{n-1}} \otimes 1 \otimes_H b_{i_1,\dots,i_{n-1};\sigma}.$$

Since C is a Var-pseudo-algebra, we have $\sum\limits_{\sigma\in S_n} b_{i_1,\dots,i_{n-1};\sigma} = 0$ for all $i_1, \dots, i_{n-1}$.

By Lemma 5

$$C_n^{(\varepsilon)}\big(t_\sigma \otimes \sigma \otimes e_j^{(n)}\big)(a_1, \dots, a_n) = \sum_{i_1,\dots,i_{n-1}} \varepsilon(h_{i_1} \overset{j}{\widehat{\dots}} h_{i_{n-1}}) h_{i_j} b_{i_1,\dots,i_{n-1};\sigma}$$

for $j \ne n$, and

$$C_n^{(\varepsilon)}(t_\sigma \otimes \sigma \otimes e_n^{(n)})(a_1, \dots, a_n) = \sum_{i_1,\dots,i_{n-1}} \varepsilon(h_{i_1} \dots h_{i_{n-1}}) b_{i_1,\dots,i_{n-1};\sigma}.$$

Therefore, $C_n^{(0)}\big(\Psi_n^{-1}\big(t \otimes e_j^{(n)}\big)\big) = C_n^{(\varepsilon)}\big(t \otimes e_j^{(n)}\big) = 0$ in $C^{(0)}(n)$, so, $C^{(0)}$ is a Var-dialgebra. ☐

It turns out that the converse is also true: a Var-dialgebra can be embedded into a pseudo-algebra (over $H = \Bbbk[T]$) that belongs to the variety Var of pseudo-algebras.

Let A be a Var-dialgebra. Since A is in particular a 0-dialgebra, there exists a pseudo-algebra C such that A is a subdialgebra of $C^{(0)}$. The purpose of this section is to show that one may choose C from the variety Var of pseudo-algebras.

Recall the construction from Proposition 4: consider $C = C_{(0)} \oplus C_{(1)}$, where $C_0 = H \otimes A$, $C_1 = (A \otimes A)/U$, $H = \Bbbk[T]$, $U = \operatorname{Span}\{\langle a,b\rangle \otimes \langle c,d\rangle \mid a,b,c,d \in A\}$, and the pseudo-product is given by (14)–(17).

As before, we identify A and C with the symmetric functors $A : \mathrm{Alg}_S \otimes \mathcal{E} \to A \in \mathrm{Vec}_\Bbbk$, $C : \mathrm{Alg}_S \to C \in H\text{-mod}$, respectively.

We will denote by $T_i^{(n)}$ the element $(T \otimes 1 \otimes \cdots \otimes 1)^{(1i)} \in H^{\otimes n}$, $i = 1, \dots, n$, and let $T_0^{(n)}$ stands for $1 \otimes \cdots \otimes 1 \in H^{\otimes n}$.

Lemma 10. *Suppose* $t \in \mathrm{Alg}(n)$, $a_1, \dots, a_n \in A$. *Then*

$$C_n(t)(a_1,\dots,a_n) = T_0^{(n)} \otimes_H x_0 - \sum_{i=1}^{n-1} T_i^{(n)} \otimes_H x_i,$$

where $x_0 \in C_{(0)}$, $x_i \in C_{(1)}$,

$$x_0 = A_n\big(t \otimes e_n^{(n)}\big)(a_1,\dots,a_n), \tag{28}$$

$$Tx_i = A_n\big(t \otimes e_n^{(n)} - t \otimes e_i^{(n)}\big)(a_1,\dots,a_n), \quad i = 1,\dots,n-1. \tag{29}$$

Proof. For $n = 2$ the statement is clear from (14). Assume we have the required relations for all terms of degree k, $k < n$. Then an arbitrary $t \in \mathrm{Alg}(n)$ can be presented as $t = \mathrm{Comp}^{(m,n-m)}(x_1x_2, t_1, t_2)$, $t_1 \in \mathrm{Alg}(m)$, $t_2 \in \mathrm{Alg}(n-m)$. By the assumption,

$$C_m(t_1)(a_1,\dots,a_m) = T_0^{(m)} \otimes_H x_0 - \sum_{i=1}^{m-1} T_i^{(m)} \otimes_H x_i,$$

$$C_{n-m}(t_2)(a_{m+1},\dots,a_n) = T_0^{(n-m)} \otimes_H y_0 - \sum_{j=1}^{n-m-1} T_j^{(n-m)} \otimes_H y_j.$$

Direct computation shows that

$$\begin{aligned} C_n(t)(a_1,\dots,a_n) &= C_m(t_1)(a_1,\dots,a_m) * C_{n-m}(t_2)(a_{m+1},\dots,a_n) \\ &= T_0^{(n)} \otimes_H (x_0 \vdash y_0) - \sum_{i=1}^{m} T_i^{(n)} \otimes_H (x_0 \otimes y_0) \\ &\quad - \sum_{i=1}^{m-1} T_i^{(n)} \otimes_H (-Tx_i) \otimes y_0 - \sum_{j=1}^{n-m-1} T_{m+j}^{(n)} \otimes_H x_0 \otimes Ty_j. \end{aligned}$$

Therefore,

$$C_n(t)(a_1,\dots,a_n) = T_0^{(n)} \otimes_H z_0 - \sum_{i=1}^{n-1} T_i^{(n)} \otimes_H z_i,$$

where

$$z_i = \begin{cases} x_0 \vdash y_0, & i = 0, \\ x_0 \otimes y_0 - Tx_i \otimes y_0, & i = 1,\dots,m-1, \\ x_0 \otimes y_0, & i = m, \\ x_0 \otimes Ty_{i-m}, & i = m+1,\dots,n-1. \end{cases}$$

By the assumption,

$$x_0 = A_m\big(t_1 \otimes e_m^{(m)}\big)(a_1,\dots,a_m), \quad y_0 = A_{n-m}\big(t_2 \otimes e_{n-m}^{(n-m)}\big)(a_{m+1},\dots,a_n).$$

Hence,

$$z_0 = A_n\big(\mathrm{Comp}^\pi\big(x_1x_2\otimes e_2^{(2)}, t_1\otimes e_m^{(m)}, t_2\otimes e_{n-m}^{(n-m)}\big)\big)(a_1,\dots,a_n) \\ = A_n\big(t\otimes e_n^{(n)}\big)(a_1,\dots,a_n),$$

where $\pi=(m,n-m)\in\Pi(n,2)$. Similarly, for $i=1,\dots,m-1$ one has

$$\begin{aligned} Tz_i &= x_0\vdash y_0 - x_0\dashv y_0 - Tx_i\vdash y_0 + Tx_i\dashv y_0 \\ &= A_n\big(\mathrm{Comp}^\pi\big(x_1x_2\otimes\big(e_2^{(2)}-e_1^{(2)}\big), t_1\otimes e_m^{(m)}, t_2\otimes e_{n-m}^{(n-m)}\big)\big)(a_1,\dots,a_n) \\ &- A_n\big(\mathrm{Comp}^\pi\big(x_1x_2\otimes\big(e_2^{(2)}-e_1^{(2)}\big), t_1\otimes\big(e_m^{(n)}-e_i^{(m)}\big), t_2\otimes e_{n-m}^{(n-m)}\big)\big)(a_1,\dots,a_n) \\ &= A_n\big(t\otimes e_n^{(n)} - t\otimes e_m^{(n)} - t\otimes e_n^{(n)} + t\otimes e_n^{(n)} + t\otimes e_m^{(n)} - t\otimes e_i^{(n)}\big)(a_1,\dots,a_n) \\ &= A_n\big(t\otimes\big(e_n^{(n)}-e_i^{(n)}\big)\big)(a_1,\dots,a_n). \end{aligned}$$

In the same way, one may proceed in the remaining cases for $i=m,m+1,\dots,n$. □

Lemma 11. *Suppose* $t\in\mathrm{Alg}_S(n)$, $t=u\otimes\sigma$, $u\in\mathrm{Alg}(n)$, $\sigma\in S_n$, *and let* $a_1,\dots,a_n\in A$. *Then*

$$C_n(t)(a_1,\dots,a_n) = T_0^{(n)}\otimes_H x_{0,\sigma} - \sum_{i=1}^{n-1} T_i^{(n)}\otimes_H x_{i,\sigma},$$

where $x_0\in C_{(0)}$, $x_i\in C_{(1)}$,

$$x_{0,\sigma} = A_n\big(t\otimes e_{n\sigma^{-1}}^{(n)}\big)(a_1,\dots,a_n), \tag{30}$$

$$x_{i,\sigma} = A_n\big(t\otimes\big(e_{n\sigma^{-1}}^{(n)} - e_{i\sigma^{-1}}^{(n)}\big)\big)(a_1,\dots,a_n),\quad i=1,\dots,n-1. \tag{31}$$

Proof. By the definition of the operad $C\in H$-mod,

$$C_n(t)(a_1,\dots,a_n) = (\sigma\otimes_H \mathrm{id})C_n(u)(a_{1\sigma},\dots,a_{n\sigma}).$$

By Lemma 10, $C_n(u)(a_{1\sigma},\dots,a_{n\sigma}) = T_0^{(n)}\otimes_H y_0 - \sum\limits_{i=1}^{n-1} T_i^{(n)}\otimes_H y_i$, where $y_0,y_1,\dots,y_n$ satisfy (28), (29).

If $n\sigma=n$ then the statement is obvious. Suppose $n\sigma\ne n$. Since $\sigma : T_i^{(n)}\mapsto T_{i\sigma}^{(n)}$, we have

$$\begin{aligned} C_n(t)(a_1,\dots,a_n) &= T_0^{(n)}\otimes_H y_0 - \sum_{i=1}^{n-1} T_{i\sigma}^{(n)}\otimes_H y_i \\ &= T_0^{(n)}\otimes_H y_0 - T_n^{(n)}\otimes_H y_{n\sigma^{-1}} - \sum_{\substack{1\le j\le n-1\\ j\ne n\sigma}} T_j^{(n)}\otimes_H y_{j\sigma^{-1}} \\ &= T_0^{(n)}\otimes_H (y_0 - Ty_{n\sigma^{-1}}) + \sum_{i=1}^{n-1} T_i^{(n)}\otimes_H (y_{n\sigma^{-1}} - (1-\delta_{i,n\sigma})y_{i\sigma^{-1}}). \end{aligned}$$

Therefore,

$$x_i = \begin{cases} y_0 - Ty_{n\sigma^{-1}}, & i=0, \\ -y_{n\sigma^{-1}} + y_{i\sigma^{-1}}, & i\ne n\sigma, \\ -y_{n\sigma^{-1}}, & i=n\sigma. \end{cases}$$

It remains to apply (28) and (29) to obtain (30) and (31). □

Lemma 12. *Suppose* $t \in \mathrm{Alg}(n)$, $a_1, \dots, a_n \in A$, $b \in A$. *Then for every* $i = 1, \dots, n$

$$C_n(t)(a_1, \dots, a_{i-1}, a_i \otimes b, a_{i+1}, \dots, a_n) = T_0^{(n)} \otimes_H x_i,$$

where $x_i \in C_{(1)}$,

$$\begin{gathered} Tx_i = A_{n+1}\big(f_i \otimes \big(e_{i+1}^{(n+1)} - e_i^{(n+1)}\big)\big)(a_1, \dots, a_i, b, a_{i+1}, \dots, a_n), \qquad (32) \\ f_i = t(x_1, \dots, x_{i-1}, x_i x_{i+1}, x_{i+2}, \dots, x_{n+1}). \end{gathered}$$

Proof. Suppose $t \in \mathrm{Alg}(n)$. If $n = 1$ then the statement is clear. Assume we are done for any $k < n$. It is sufficient to show the required relations for $t = (x_1 \dots x_n) \in \mathrm{Alg}(n)$. Let us present t as $t = \mathrm{Comp}^{(m,n-m)}(x_1x_2, t_1, t_2)$, $1 \le m \le n$, $t_1 = (x_1 \dots x_m)_1$, $t_2 = (x_1 \dots x_{n-m})_2$, where $(\dots)_1$, $(\dots)_2$ are some bracketings. Suppose $i \in \{1, \dots, m\}$. Then

$$\begin{aligned} &C_n(t)(a_1, \dots, a_i \otimes b_i, \dots a_n) \\ &\qquad = C_m(t_1)(a_1, \dots, a_i \otimes b_i, \dots a_m) * C_{n-m}(t_2)(a_{m+1}, \dots, a_n). \end{aligned}$$

By the assumption

$$C_m(t_1)(a_1, \dots, a_i \otimes b, \dots, a_m) = T_0^{(m)} \otimes_H x_i,$$

where $x_i \in C_{(1)}$ satisfies (32), and by Lemma 10

$$C_{n-m}(t_2)(a_{m+1}, \dots, a_n) = T_0^{(n-m)} \otimes_H y_0 - \sum_{j=1}^{n-m-1} T_j^{(n-m)} \otimes_H y_j,$$

$y_0 = A_{n-m}\big(t_2 \otimes e_{n-m}^{(n-m)}\big)(a_{m+1}, \dots, a_n)$, $y_j \in C_{(1)}$ for $j = 1, \dots, n-m$.

Since $C_{(1)} * C_{(1)} = 0$, we have

$$C_n(t)(a_1, \dots, a_i \otimes b, \dots, a_n) = \big(T_0^{(m)} \otimes_H x_i\big) * \big(T_0^{(n-m)} \otimes_H y_0\big) = -T_0^{(n)} \otimes_H (Tx_i \otimes y_0).$$

Hence, it is enough to consider $-T(Tx_i \otimes y_0) = Tx_i \dashv y_0 - Tx_i \vdash y_0$. Recall that

$$Tx_i = A_{m+1}\big(f_i \otimes \big(e_{i+1}^{(m+1)} - e_i^{(m+1)}\big)\big)(a_1, \dots, a_i, b, \dots, a_m),$$

where $f_i = t_1(x_1, \dots, x_i x_{i+1}, \dots, x_{m+1})$. Therefore,

$$Tx_i \dashv y_0 - Tx_i \vdash y_0 = F_i(a_1, \dots, a_i, b, \dots, a_n),$$

where

$$\begin{aligned} F_i &= \mathrm{Comp}^{(m+1,n-m)}\big(x_1x_2 \otimes \big(e_1^{(2)} - e_2^{(2)}\big), f_i \otimes \big(e_{i+1}^{(m+1)} - e_i^{(m+1)}\big), t_2 \otimes e_{n-m}^{(n-m)}\big) \\ &= t_1(x_1, \dots, x_i x_{i+1}, \dots x_{m+1}) t_2(x_{m+2}, \dots, x_n) \otimes \big(e_i^{(n+1)} - e_i^{(n+1)}\big) \\ &= t(x_1, \dots, x_i x_{i+1}, \dots x_{n+1}) \otimes \big(e_i^{(n+1)} - e_i^{(n+1)}\big). \end{aligned}$$

The remaining case $i \in \{m+1, \dots, n\}$ is completely analogous. □

Lemma 13. *Suppose* $t \in \mathrm{Alg}_S(n)$, $t = u \otimes \sigma$, $u \in \mathrm{Alg}(n)$, $\sigma \in S_n$, *and let* $a_1, \dots, a_n \in A$, $b \in A$. *Then for every* $i = 1, \dots, n$

$$C_n(t)(a_1, \dots, a_{i-1}, a_i \otimes b, a_{i+1}, \dots, a_n) = T_0^{(n)} \otimes_H x_i,$$

where $x_i \in C_{(1)}$,

$$\begin{gathered} Tx_i = A_{n+1}\big(f_i \otimes \big(e_{i\sigma^{-1}+1}^{(n+1)} - e_{i\sigma^{-1}}^{(n+1)}\big)\big)(a_1, \dots, a_i, b, a_{i+1}, \dots, a_n), \qquad (33) \\ f_i = t(x_1, \dots, x_{i-1}, x_i x_{i+1}, x_{i+2}, \dots, x_{n+1}). \end{gathered}$$

Proof. By the definition of the operad $C \in H$-mod,

$$C_n(t)(a_1,\dots,a_{i-1},a_i\otimes b,a_{i+1},\dots,a_n) = (\sigma\otimes_H \mathrm{id})C_n(u)(z_1,\dots,z_n),$$

where $z_j = a_{j\sigma}$ for $j \neq i\sigma^{-1}$, and $z_{i\sigma^{-1}} = a_i\otimes b$. By Lemma 12 we get the required relation. □

Theorem 14. *For every* Var*-dialgebra A there exists a* Var*-pseudo-algebra $C_{\mathrm{Var}}(A)$ such that $A\subseteq C_{\mathrm{Var}}(A)^{(0)}$.*

Proof. Consider the pseudo-algebra $C = C(A)$ from Proposition 4. It is enough to show that there exists an ideal I of C such that $I\cap C_{(0)} = 0$ and C/I is a Var-pseudo-algebra.

Recall that Σ is the system of polylinear defining identities of the variety Var. For any $t\in\Sigma$ of degree $n\geq 2$, consider the expressions

$$C_n(t)(a_1,\dots,a_n) = T_0^{(n)}\otimes_H x_0 - \sum_{i=1}^{n-1} T_i^{(n)}\otimes_H x_i, \tag{34}$$

$$C_n(t)(a_1,\dots,a_{j-1},a_j\otimes b,a_{j+1},\dots,a_n) = T_0^{(n)}\otimes_H y_j, \tag{35}$$

for all $a_1,\dots,a_n,b\in A$, $j=1,\dots,n$. By (28) and (30) we have $x_0=0$.

Denote by I the linear span of all elements $x_i, y_j\in C_{(1)}$ from (34), (35), $i,j\geq 1$, for all $a_1,\dots,a_n,b\in A$ and for all $t\in\Sigma$. It follows from Lemmas 10–13 and (15), (16) that I is actually an ideal of the pseudo-algebra C. Moreover, it is clear that $C_{\mathrm{Var}}(A) = C/I$ (as a pseudo-algebra) satisfies all of the identities $t\in\Sigma$: if two or more arguments among $x_1,\dots,x_n\in C$ belong to $C_{(1)}$ then $C_n(t)(x_1,\dots,x_n)=0$ by (17). □

Remark 15. Theorem 14 provides another proof of Proposition 7.

Indeed, if A is a 0-dialgebra that satisfies $t\otimes e_i^{(n)}$, $t\in\Sigma$, $\deg t = n$, $i=1,\dots,n$, then there exists a Var-pseudo-algebra C such that $A\subset C^{(0)}$. If $g\in \mathrm{Alg}_S(n)$, $\mathrm{Var}_n(g)=0$, then by Lemma 5 we have $C_n^{(\varepsilon)}\big(g\otimes e_i^{(n)}\big)=0$ for all $i=1,\dots,n$. Therefore, $g\otimes e_i^{(n)}$ is an identity on $C^{(0)}\supset A$.

Proposition 16. *Let A be a* Var*-dialgebra, and let P be a* Var*-pseudo-algebra over $H=\Bbbk[T]$ such that there exists a homomorphism of dialgebras $\varphi: A\to P^{(0)}$ and $\varphi(a)*\varphi(b) = \sum\limits_i h_i\otimes 1\otimes_H c_i$, $\deg h_i\leq 1$ for all $a,b\in A$. Then there exists a homomorphism of pseudo-algebras $C_{\mathrm{Var}}(\varphi): C_{\mathrm{Var}}(A)\to P$ such that $C_{\mathrm{Var}}(\varphi)|_A = \varphi$.*

Proof. Let us first consider the case when $\Sigma=\emptyset$, i.e., A is an arbitrary 0-dialgebra.

Suppose P is a pseudo-algebra over $H=\Bbbk[T]$ that satisfies the conditions of the proposition. Since $\Bbbk$ may be of positive characteristic, we need a slightly different definition of n-products:

$$x_{\ (n)}\ y = z_n,\quad x,y\in P,$$

where

$$x*y = \sum_{s\geq 0} T^s\otimes 1\otimes_H z_s.$$

Then $x_{\ (n)}\ y = 0$ for $n\gg 0$, and $H^{\otimes 2}$-linearity implies

$$Tx_{\ (n)}\ y = x_{\ (n-1)}\ y,\quad x_{\ (n)}\ Ty = T(x_{\ (n)}\ y) - x_{\ (n-1)}\ y, \tag{36}$$

assuming $x \,{}_{(-1)}\, y = 0$.

For all $a, b \in A$ we have

$$\varphi(a \vdash b) = \varphi(a) \,{}_{(0)}\, \varphi(b), \quad \varphi(a \dashv b) = \varphi(a) \,{}_{(0)}\, \varphi(b) + T(\varphi(a) \,{}_{(1)}\, \varphi(b)) \tag{37}$$

since $\varphi : A \to P^{(0)}$ is a homomorphism of dialgebras.

Lemma 17. *For all $a, b, c, d \in A$ we have $(\varphi(a) \,{}_{(1)}\, \varphi(b)) \,{}_{(n)}\, (\varphi(c) \,{}_{(1)}\, \varphi(d)) = 0$ for $n \geq 0$, and $(\varphi(a) \,{}_{(1)}\, \varphi(b)) \,{}_{(n)}\, \varphi(c) = 0$, $\varphi(a) \,{}_{(n)}\, (\varphi(b) \,{}_{(1)}\, \varphi(c)) = 0$ for $n \geq 1$.*

Proof. Denote $\varphi(a)$, $\varphi(b)$, $\varphi(c)$, $\varphi(d)$ by x, y, z, u, respectively.

There exists $N \geq 0$ such that $x \,{}_{(n)}\, (y \,{}_{(1)}\, z) = 0$ for all $n \geq N$. Suppose that the minimal of such N greater than 2. Then by (37)

$$x \,{}_{(N)}\, T(y \,{}_{(1)}\, z) = x \,{}_{(N)}\, \varphi(b \vdash c) - x \,{}_{(N)}\, \varphi(b \dashv c) = 0.$$

On the other hand, by (36)

$$x \,{}_{(N)}\, T(y \,{}_{(1)}\, z) = T(x \,{}_{(N)}\, (y \,{}_{(1)}\, z)) - x \,{}_{(N-1)}\, (y \,{}_{(1)}\, z),$$

so $x \,{}_{(N-1)}\, (y \,{}_{(1)}\, z) = 0$ in contradiction with the minimality of N. Other statements of the Lemma can be proved in a completely similar way. □

Define a map $\psi : C(A) \to P$ by the following rule:

$$\psi(h \otimes a) = h(T)\varphi(a), \quad \psi(a \otimes b) = -\varphi(a) \,{}_{(1)}\, \varphi(b).$$

The second equation actually means that we define $\psi : A \otimes A \to P$, then by Lemma 17 we have $U = \operatorname{Span}\{\langle a, b\rangle \otimes \langle c, d\rangle \mid a, b, c, d \in A\} \subseteq \operatorname{Ker} \psi$.

It remains to show that ψ is actually a homomorphism of pseudo-algebras. Relation (37) implies that ψ is H-linear. To check that ψ preserves $*$, one should proceed with four cases following (14)–(17). Let us consider the second case for example: $a * (b \otimes c) = 1 \otimes 1 \otimes_H a \otimes \langle b, c\rangle$, $\psi(a \otimes \langle b, c\rangle) = -\varphi(a) \,{}_{(1)}\, \varphi(\langle b, c\rangle) = \varphi(a) \,{}_{(1)}\, T(\varphi(b) \,{}_{(1)}\, \varphi(c)) = T(\varphi(a) \,{}_{(1)}\, (\varphi(b) \,{}_{(1)}\, \varphi(c))) - \varphi(a) \,{}_{(0)}\, (\varphi(b) \,{}_{(1)}\, \varphi(c)) = -\varphi(a) \,{}_{(0)}\, (\varphi(b) \,{}_{(1)}\, \varphi(c))$ by Lemma 17. On the other hand, $\psi(a) * \psi(b \otimes c) = -1 \otimes 1 \otimes_H \varphi(a) \,{}_{(0)}\, (\varphi(b) \,{}_{(1)}\, \varphi(c))$.

Now, let Σ be a nonempty family of polylinear identities defining a homogeneous variety Var, and let A be a Var-dialgebra. As we have proved, for any pseudo-algebra P satisfying the conditions of the theorem there exists a homomorphism of pseudo-algebras $C(\varphi) : C(A) \to P$ such that $C(\varphi)|_A = \varphi$. Recall that $C_{\mathrm{Var}}(A) = C/I$, where I is spanned by all coefficients of $C(A)_n(t^*)(u_1, \dots, u_n) \in H^{\otimes n} \otimes_H C(A)$, $t \in \Sigma$, $n = \deg t$, $u_i \in C(A)$, $i = 1, \dots, n$. Therefore, $I \subseteq \operatorname{Ker} C(\varphi)$ and there exists a homomorphism $C_{\mathrm{Var}}(\varphi)$ with the required properties. □

5. Conformal representations of Leibniz algebras

Recall the notions of a conformal endomorphism [6, 11] and a representation of an (associative or Lie) pseudo-algebra [9]. We will generally follow [10] in the exposition.

Consider a left module M over $H = \Bbbk[T]$. By $\operatorname{End} M$ we denote the associative algebra of $\Bbbk$-linear maps from M to itself. A map $a : \Bbbk \to \operatorname{End} M$ is said to be *locally regular* if for any $u \in M$ the map $z \mapsto a(z)u$ is a regular M-valued map on the affine line $\Bbbk$.

Note that $\operatorname{End} M$ can be considered as a left H-module with respect to

$$(f\varphi)(u) = f_{(2)}\varphi(f_{(-1)}u), \quad f \in H,\ u \in M,\ \varphi \in \operatorname{End} M.$$

A map $a : \Bbbk \to \operatorname{End} M$ is *translation-invariant* if

$$fa(z) = f(z)a(z), \quad f \in A,\ z \in V.$$

Definition 3. A *conformal endomorphism* of an H-module M is a locally regular and translation invariant map $a : \Bbbk \to \operatorname{End} M$.

The set $\operatorname{Cend} M$ of all conformal endomorphisms of M is a linear space which is also an H-module with respect to the action defined by

$$(fa)(z) = f(z)a(z), \quad f \in H,\ a \in \operatorname{Cend} M,\ z \in \Bbbk.$$

It is easy to see that an H-pseudo-algebra structure on a left H-module C can be equivalently defined via an H-linear map $\mu : C \to \operatorname{Cend} C$. Indeed, the correspondence between the map μ and pseudo-product $*$ is completely described as follows: if

$$\mu(a)(x)b = \sum_{s=0}^{n} x^s c_s, \quad b, c_s \in C,$$

then

$$a * b = \sum_{s=0}^{n} (-T)^s \otimes 1 \otimes_H c_s.$$

For the H-module $\operatorname{Cend} M$ and for every $a \in \operatorname{Cend} M$ one may define a map $(a_x\cdot) : \Bbbk \to \operatorname{End}\operatorname{Cend} M$ as follows:

$$(a_z b)(w) = a(z)b(w - z), \quad b \in \operatorname{Cend} M,\ z, w \in \Bbbk.$$

This map may not be locally regular, so $\operatorname{Cend} M$ is not necessarily a pseudo-algebra (see [9, 6, 11]). However, some of its submodules are actually pseudo-algebras. For example, if M is a free H-module generated by a space M_0 (i.e., $M = H \otimes M_0$) then the current submodule $\operatorname{Cur}\operatorname{End} M_0 = H \otimes \operatorname{End} M_0 \subset \operatorname{Cend} M$, where $(f \otimes \varphi)(z) = f(-z)(\mathrm{id}_H \otimes \varphi)$, $f \in H$, $\varphi \in \operatorname{End} M_0$, $z \in \Bbbk$, is always an associative pseudo-algebra.

Every associative pseudo-algebra C can be considered as a Lie pseudo-algebra with respect to new pseudo-product

$$[a * b] = a * b - ((12) \otimes_H \mathrm{id}_C)(b * a).$$

The Lie pseudo-algebra obtained is denoted by $C^{(-)}$. In particular, if the entire $\operatorname{Cend} M$ is a pseudo-algebra (that is the case if M is a finitely generated H-module) then $\operatorname{Cend} M^{(-)}$ is denoted by $\operatorname{gc} M$. Even if $\operatorname{Cend} M$ is not a pseudo-algebra, we will use the following convention: $L \subset \operatorname{gc} M$ means that L is a subalgebra of some $C^{(-)}$, where $C \subseteq \operatorname{Cend} M$ is a pseudo-algebra.

Definition 4. Let $\mathfrak{g}$ be a Leibniz algebra, i.e., a Lie dialgebra. A homomorphism of dialgebras $\rho : \mathfrak{g} \to L^{(0)}$, where $L \subseteq \operatorname{gc} M$, is called a *conformal representation* of $\mathfrak{g}$ on M.

A conformal representation is said to be finite if M is a finitely generated H-module.

Proposition 18. *Every Leibniz algebra $\mathfrak{g}$ has a faithful conformal representation. If $\dim \mathfrak{g} < \infty$ then there exists a finite faithful conformal representation.*

Proof. Let us use $[ab]$ for $a \dashv b$, $a, b \in \mathfrak{g}$. Denote by $\mathfrak{l}$ the (ordinary) Lie algebra $\mathfrak{g}/\operatorname{Span}\{[xx] \mid x \in \mathfrak{g}\}$. If $x \in \mathfrak{g}$ then $\bar{x}$ is the image of x in $\mathfrak{l}$.

Let V stands for an arbitrary nonzero (left) $\mathfrak{l}$-module. Set $M_0 = V \oplus (\mathfrak{g} \otimes V)$, $M = H \otimes M_0$. Consider the map

$$\rho : \mathfrak{g} \to \operatorname{Cend} M$$

defined as follows:

$$\begin{aligned} \rho(x) * u &= 1 \otimes 1 \otimes_H \bar{x}u - T \otimes 1 \otimes_H (x \otimes u), \\ \rho(x) * (a \otimes u) &= 1 \otimes 1 \otimes_H (a \otimes \bar{x}u - [ax] \otimes u), \end{aligned} \tag{38}$$

$a, x \in \mathfrak{g}$, $u \in V$. It is easy to see that $\rho(x) \in \operatorname{Cur}\operatorname{End} M_0$. Indeed,

$$\rho(x) = \rho_0(x) - T\rho_1(x),$$

where $\rho_i(x) \in \operatorname{End} M_0$, $\rho_0(x) = \bar{x} \oplus (\operatorname{id} \otimes \bar{x} - [\cdot x] \otimes \operatorname{id})$, $\rho_1(x)$ maps $\mathfrak{g} \otimes V$ to zero and $u \in V$ to $x \otimes u$. Therefore, ρ is injective, and $\rho(\mathfrak{g})$ generates an associative pseudo-algebra C that lies in $\operatorname{Cur}\operatorname{End} M_0$. Moreover,

$$\begin{aligned} &\rho(a) * \rho(b) = (\rho_0(a) - T\rho_1(a)) * (\rho_0(b) - T\rho_1(b)) \\ &= 1 \otimes 1 \otimes_H \rho_0(a)\rho_0(b) - T \otimes 1 \otimes_H \rho_1(a)\rho_0(b) - 1 \otimes T \otimes_H \rho_0(a)\rho_1(b) + T \otimes T \otimes_H \rho_1(a)\rho_1(b) \\ &= 1 \otimes 1 \otimes_H (\rho_0(a)\rho_0(b) - T\rho_0(a)\rho_1(b)) - T \otimes 1 \otimes_H (\rho_1(a)\rho_0(b) - \rho_0(a)\rho_1(b)), \end{aligned} \tag{39}$$

since $\rho_1(a)\rho_1(b) = 0$.

This is straightforward to check that

$$[\rho_0(a), \rho_0(b)] = \rho_0([ab]), \quad [\rho_1(a), \rho_0(b)] = \rho_1([ab]).$$

Then (12) and (39) imply

$$C_2^{(0)}(x_1 \dashv x_2 - x_2 \vdash x_1)(\rho(a), \rho(b)) = \rho([ab]), \quad a, b \in \mathfrak{g},$$

that means ρ is a conformal representation of $\mathfrak{g}$.

If $\mathfrak{g}$ is finite-dimensional then so can be choosen V, and hence M is a finitely generated H-module. □

Corollary 19 ([3]). *Every Leibniz algebra can be embedded into an associative dialgebra.*

Proof. Consider a Leibniz algebra $\mathfrak{g}$ and its faithful conformal representation $\rho : \mathfrak{g} \to L \subset \operatorname{gc} M$ as in Proposition 18. We have seen that L is actually a Lie subalgebra of $(\operatorname{Cur}\operatorname{End} M_0)^{(-)}$, so $\mathfrak{g} \subset L \subset A = (\operatorname{Cur}\operatorname{End} M_0)^{(0)}$. The latter is an associative dialgebra by Theorem 9. □

Remark 20. If $V = U(\mathfrak{l})$ in Proposition 18, then the associative dialgebra generated in $(\operatorname{Cend} M)^{(0)}$ by the image of $\mathfrak{g}$ is isomorphic to the universal enveloping dialgebra $Ud(\mathfrak{g})$ constructed in [3]. However, $C_{Lie}(\mathfrak{g})$ is not embedded into $\operatorname{Cur}\operatorname{End} M_0$ for $M_0 = U(\mathfrak{l}) \oplus (\mathfrak{g} \otimes U(\mathfrak{l}))$.

REFERENCES

[1] J.-L. Loday, Une version non commutative des alg‘ebres de Lie: les alg‘ebres de Leibniz, Enseign. Math., 39 (1993) 269-293.

[2] J.-L. Loday, T. Pirashvili, Universal envelopping algebras of Leibniz algebras and homology, Math. Ann. **296** (1993) 139–158.

[3] J.-L. Loday, Dialgebras, in: Dialgebras and Related Operads, Lecture Notes in Mathematics, vol. 1763, pp. 7-66. Springer Verl., Berlin, 2001.

[4] D. Liu, Steinberg–Leibniz algebras and superalgebras, J. Algebra 283 (1) (2005) 199–221.

[5] J.R. Faulkner, Barbilian planes, Geom. Dedicata 30 (1989) 125–181.

[6] V.G. Kac, Vertex Algebras for Beginners, University Lecture Series, vol. 10, AMS, Providence, RI, 1996.
[7] M. Roitman, On free conformal and vertex algebras, J. Algebra 217 (2) (1999) 496–527.
[8] P.S. Kolesnikov, Identities of conformal algebras and pseudoalgebras, Comm. Algebra 34 (6) (2006) 1965–1979.
[9] B. Bakalov, A. D'Andrea, V.G. Kac, Theory of finite pseudoalgebras, Adv. Math. 162 (1) (2001) 1–140.
[10] P.S. Kolesnikov, Associative algebras related to conformal algebras, Applied Categorical Structures, to appear.
[11] V.G. Kac, Formal distribution algebras and conformal algebras, XII-th International Congress in Mathematical Physics (ICMP'97) (Brisbane), International Press, Cambridge, MA, 1999, pp. 80–97.

E-mail address: `pavelsk@math.nsc.ru`

Sobolev Institute of Mathematics, Novosibirsk, Russia